

Medical sciences

УДК 618.4

Yalonetskiy I.Z.

assistant

Belarusian State Medical University

Zhurova A.V.

student

Belarusian State Medical University

Serbina D.V.

student

Belarusian State Medical University

QUALITY OF LABOR ANALGESIA DEPENDING ON ANESTHETIC TACTICS

Summary: *The comparison of spinal analgesia and obstetric sleep and their effect on delivery process and the baby.*

Key words: labor analgesia, obstetric sleep, spinal analgesia.

The problem of pain relief in labor is not only medical but also social. Often after painful labor woman acquires fear of further childbearing, which negatively affects the country's demographics. Besides, the risk of postnatal depression increases. [1, P. 7-11] Therefore, ensuring a comfortable childbirth should be a priority task for obstetric and anesthetic services.

Stress reactions of maternal body can also be successfully corrected or even eliminated with adequate analgesia [2, P. 150-156].

It is also necessary to pay attention to how labor anesthesia may affect the fetus. According to studies, neuraxial blockade does not increase the duration of labor and does not affect the fetus, which is important to improve the level of public health [3, P. 38-50; 4, P. 6-10]. Currently in the US, this method is used

in more than 60% of labors. In Belarus its using is less than 20% of cases. [5, P. 28-31]

Obstetric sleep continues to be applied in our country which generates controversy between anesthesiologists and obstetricians because the feasibility of the method is considered doubtful among anesthesiologists.

In order to understand this issue, we compared the quality of labor analgesia with spinal analgesia with low doses of bupivacaine and obstetric sleep and the impact of these methods on the mother and fetus.

We examined 45 patients of 6th and 5th Minsk City Clinical Hospital. The study included mothers who agreed to participate in age from 20 to 35 years, gestational age from 255 to 285 days, delivered per vias naturales and who had no comorbidity.

New mothers were divided into 3 equal groups of 15 people. Group A (control) consisted of patients who gave birth without anesthesia. Group B included 15 women who gave birth with spinal analgesia (SA). Blockade was performed at the level L_{III-IV} with MarkainSpinalHevi (hyperbaric bupivacaine solution) in a dose of 1.5 mg. Group C included women with obstetric sleep (MS) with using of Hydroxybutyric acid (GHB) in a dose of 50 mg / kg, which lasted 1.5 hours \pm 18,25 min.

Groups did not differ in age, height, maternal weight, gestational period and fetal weight.

Duration of delivery by group is shown in Table 1 and averaged 7 hours \pm 20 min 2 hours. No statistically significant differences were detected between the groups. Thus, the choice of anesthetic tactics did not have a significant effect on the duration of labor.

Total duration of labor

	Group A	Group B	Group C
Length, m	420,33 [330; 510]	467 [380; 460]	475,33 [400; 550]

Table 1. authors' design.

Hemodynamics of mothers during all periods of childbirth was assessed by arterial pressure and heart rate, but the difference between both groups and within groups was not statistically significant.

Severity of pain was assessed using a visual analogue scale (VAS) [6, P. 7-12] and numeral rating scale (NRS) [7, P. 697-706]. VAS is a straight line length of 10 cm. The patient was asked to make a mark on the line corresponding to the intensity of his pain. Starting point line represents no pain - 0, then there is a weak, moderate, strong and unbearable pain - 10. Digital rating scale consists of 11 items 0 "no pain" and 10 "unbearable pain".




Figure 1. - VAS and NRS, authors' design.

Patients of the groups A and C suffered from severe pain, while a patient from Group B, which was held spinal analgesia, observed more comfortable labor, estimating maximum pain at 6 points out of 10 in the second period. The difference between the two groups, calculated by the Mann-Whitney test, was statistically significant in the first and second stage of labor ($p < 0,05$), and the index difference inside the groups in all periods ($p < 0,05$ Student's criterion). The results are presented graphically in figure 2.




Figure 2. – Severity of pain, authors’ design.

Fetal status was assessed according to the CTG, has been based upon the basal heart rate and STV-rhythm (ShortTermVariations) as indicators of fetal hypoxia. However significant differences between the groups haven’t been identified, from which it can be concluded that the anesthetic did not have a pronounced effect on the fetus. In assessing newborns on Apgar score at 1 and 5 minutes of life the difference between groups A and B was not significant, but the group C figures were significantly lower ($p < 0,05$; $U \geq 2$).




Figure 3. – Newborns’ indicators, authors’ design.

Also in Group C in 5 cases labors resulted in caesarean section; were revealed a weakening of labor (n = 4), signs of fetal hypoxia (n = 3); fetal depression (n = 1).

Summary, we can make following conclusions. First, spinal anesthesia with low doses of bupivacaine promotes comfortable labor without providing negative impact on woman and fetus and don't increase risk of complications. Second, use of obstetric sleep may cause the weakening of labor and fetal depression.

Literature:

1. Кобеляцкий, Ю.Ю. Обезболивание родов в условиях новой акушерской доктрины / Ю. Ю. Кобеляцкий, Л. В. Ефремова // Острые и неотложные состояния в практике врача. – 2009. - №4. – С. 7-11.
2. Эйзенах Дж. Родовая боль / Дж. Эйзенах // Актуальные проблемы анестезиологии и реаниматологии. Освежающий курс лекций. – 2007. - №12. – С. 150-156.
3. Reynolds F. Labour analgesia and the baby: good news is no news / F. Reynolds // Int. J. of Obst. Anesthesia -. 2011. - Vol. 20. - P. 38-50.
4. Шифман Е. М. Влияние эпидуральной анальгезии на течение родов / Е. М. Шифман, Ю. Н. Ермилов // Рос. вестн. акушера-гинеколога. – 2006. - №2. – С. 6-10.
5. Малюк В.В. Акушерские и анестезиологические особенности ведения родов на фоне продленной эпидуральной анальгезии / В.В. Малюк, В.В. Базовкин // Жіночий лікар. – 2010. - №6. – С. 28-31.
6. Bijur P. Reliability of the visual analog scale for measurement of acute pain / PE Bijur, W. Silver, EJ Gallagher // Academic Emergency Medicine -. 2001. - № 8. - P. 7-12.
7. Gould, D. Informational point: Visual Analogue Scale (VAS) / D. Gould // Journal of Clinical Nursing - 2001. - № 10. - P. 697-706.