

Medical sciences

UDC 616.71-001.5

**Minkov Dimitar**

*MD PhD*

*Medical University Pleven,*

*Institute of Science and Research,*

*Avis Medica Hospital, Department of Orthopedics and Traumatology*

**Evtimov Kalin**

*Avis Medica Hospital, Department of Orthopedics and Traumatology*

## **FRACTURE OF HAMATE BONE: CASE REPORT**

**Summary.** *Fractures of the hamate bone are quite rare. The presented clinical case refers to a patient with a type 2b fracture according to the classification of Hirano and Inoue.*

**Key words:** *hamate bone, hamate fracture.*

**Abbreviation:** ORIF – open reposition and internal fixation; K- wires – Kirschner wires.

### **Introduction**

Fractures of the hamate bone are quite rare, constituting about 2% of all carpal fractures [1, 2]. However, two classifications have been created for them. Milch's classification distributes fractures of hamate bone in two main types of fractures: type 1 – fracture of the hook of the hamate and type 2 – fracture of the body of the hamate. *Hirano and Inoue* further refine the Milch's classification as sub-divided body fractures into type 2 coronal fractures of the body (with subtypes 2a dorsal oblique fractures and 2b splitting fractures), and type 3 – a transverse fracture of the body [3]. Moreover, they differentiated each type 2a fracture as a dorsal oblique or a splitting fracture (Fig. 1).




Fig. 1. Classification of hamate fractures (after Hirano and Inoue, 2005)

### Case report

We report the case of a 19-year-old male, who injured his right wrist during street brawl. Initial X-ray films of the right hand was made in Emergency Center 12 hours after the trauma demonstrated a type 2b fracture of the hamate (Fig. 2).



Fig. 2. X-ray images of right wrist

A computer tomography study with a 3D reconstruction was performed. The results are represented on Fig. 3.



**F**

**Fig. 3. CT-scan and 3D reconstruction of the fracture of hamate bone**

An operative treatment was performed and was inserted metal osteosynthesis K-wires. Treatment continued with cast immobilization for 35 days (Fig. 4).



**Fig. 4. Postoperative X-ray images of right wrist**

## Discussion

The type 1 fractures are often the result of repeated micro trauma to the hook during sports involving swinging clubs, bats, or racquets and direct trauma

too. *Kapickis and al.* even suggest that their frequency will grow as the popularity of rocket sports and golf increases [4]. The mechanism of fracture in type 2 fractures is direct trauma.

*Hirano and Inoue* describe 11 fractures of the type 2a all of them were occurred with a clenched fist against a wall or floor. Five cases were treated with closed reduction and percutaneous pinning and the remaining six received ORIF with screws or K-wires [3]. In our case was performed a surgical treatment ORIF with K-wires. Coronal fractures of the body of the hamate are frequently associated with ulnar CMC joint dislocations [5]. In our case was not established CMC dislocation.

### **References**

1. Rockwood and Green's fractures in adults. 6th ed. Philadelphia, Pa: Lippincott Williams & Wilkins, 2006.
2. Lister G. The Hand: Diagnosis and Indications. 3rd ed. Philadelphia, Pa: Churchill Livingstone; 1993. - p. 88–92.
3. Hirano K, Inoue G. Classification and treatment of hamate fractures. *Hand Surg.* 2005;10(2-3):151-7.
4. Kapickis M, Looi KP, Chong AK. Combined fractures of the body and hook of hamate: A form of ulnar axial injury of the wrist. *Scand J Plast Reconstr Surg Hand Surg*, 2005; 39: 116/119.
5. Cain JE, Thomas R, Wilson MR, Hematometacarpal fracture-dislocation: classification and treatment, *JHandSurg* 12A:762–767, 1987.